



CrackingCHEM

GCSE | A-LEVEL

UNDERSTAND • APPLY • SUCCEED

OCR A H432

Bonding I

Original exam questions | *TEST IT* → *EXAM STYLE* → *HARDER MIXED* → *STRETCH*

SPECIFICATION-LED. PROGRESSIVE. EXAM-FOCUSED.

OCR A SPECIFICATION

2.1.2(a) | 2.2.2(a-f,n-o) | linked 3.1.1(d-f)

TEST IT	EXAM STYLE	HARDER MIXED	STRETCH
8	10	5	1

24 ORIGINAL QUESTIONS

166 TOTAL MARKS

FINAL NOTES-ALIGNED EDITION | FORMULAE • BONDING • STRUCTURE • PROPERTIES

Independent practice material. Every question is original and stays within the finished OCR A Bonding I notes.

**START HERE**

How to use this pack

Define precisely, prove electron origin, then build every property explanation as a chain

- 1 Complete TEST IT without the notes. Correct weak definitions before moving on.
- 2 For every diagram, check electron origin, electron totals, brackets and charges.
- 3 For every property, write structure → particles → attraction → energy or mobility → property.
- 4 Use the marks to count distinct points. Do not replace precise particles with vague words such as things or particles.

✓ **OCR A boundary:** This pack covers 2.1.2(a), 2.2.2(a-f,n-o) and only the Bonding I foundation links in 3.1.1(d-f). Shapes, bond angles, electronegativity, polarity, detailed intermolecular forces and the full silicon/graphene Periodicity treatment are excluded.

Definition discipline: Electron transfer forms ions. The ionic bond is the electrostatic attraction between the resulting oppositely charged ions. A covalent definition must name the shared pair and both nuclei.

Diagram standard: Use dots and crosses consistently. Ionic diagrams require square brackets and charges. Dative arrows start at the donated electron pair and point to the acceptor.

Property route: Start with the structure, name the particles and attraction, then state the energy needed or whether charged particles are mobile.

Notes alignment: All questions map directly to the final CrackingChem OCR A Bonding I notes and the current OCR A H432 specification.

THE FIVE-LINK PROPERTY ROUTE

STRUCTURE	PARTICLES	ATTRACTION	ENERGY / MOBILITY	PROPERTY
giant or simple	ions, atoms, molecules or electrons	name it precisely	overcome it or move charge	state the observed result



SECTION A

TEST IT

Eight direct checks of the finished-notes mastery outcomes

QUESTION 1 | TEST IT

Complete the table of OCR A ions.

Ion name	Formula and charge
nitrate	
carbonate	
sulfate	
hydroxide	
ammonium	
zinc	
silver	

[7 marks]

QUESTION 2 | TEST IT

Write the formula of each compound. Every formula must be neutral and in the smallest whole-number ratio.

Compound	Formula [6 marks]
aluminium oxide	
calcium nitrate	
ammonium sulfate	
magnesium hydroxide	
zinc carbonate	
silver nitrate	

(g) State when brackets are required around a compound ion.

[1 mark]

**QUESTION 3** | **TEST IT**

(a) State the precise OCR A definition of ionic bonding.

[2 marks]

(b) Explain why electron transfer is not itself the ionic bond.

[2 marks]

QUESTION 4 | **TEST IT**

Draw a dot-and-cross diagram for magnesium chloride, MgCl_2 . Show outer-shell electrons only, electron origin, square brackets, charges and complete outer shells.

DIAGRAM SPACE

[5 marks]

**QUESTION 5** | **TEST IT**

(a) State the precise OCR A definition of a covalent bond.

[3 marks]

(b) State the number of shared electron pairs in H_2 , O_2 and N_2 .

[2 marks]

QUESTION 6 | **TEST IT**

Draw complete dot-and-cross diagrams for HCl , O_2 and N_2 . Show all shared pairs and all lone pairs.

THREE MODEL DIAGRAMS

[7 marks]



(a) State what is meant by a dative covalent bond.

[2 marks]

(b) For $\text{NH}_3 + \text{H}^+ \rightarrow \text{NH}_4^+$, name the electron-pair donor and acceptor and state the arrow direction.

[2 marks]

(c) Draw the ammonium ion, showing the origin of the electron pair in the newly formed bond, square brackets and overall charge.

DIAGRAM SPACE

[2 marks]

(a) State the precise definition of metallic bonding.

[2 marks]

(b) Bond X has an average bond enthalpy of 436 kJ mol^{-1} and bond Y has 598 kJ mol^{-1} . Identify the stronger bond and justify your choice.

[1 mark]



SECTION B

EXAM STYLE

Ten original structured questions using OCR A bonding and structure conventions

✓ **Route check:** Name the structure, particles and attraction before discussing energy or mobility.

QUESTION 9 | EXAM STYLE

Sodium chloride forms a giant ionic lattice.

(a) Explain why sodium chloride has a high melting point.

.....

.....

.....

[2 marks]

(b) Explain why solid sodium chloride does not conduct electricity but molten sodium chloride does.

.....

.....

.....

.....

[3 marks]

(c) Explain why aqueous sodium chloride conducts electricity.

.....

.....

.....

[1 mark]

(d) Write one careful sentence about the solubility of ionic compounds in water.

.....

.....

.....

[1 mark]

**QUESTION 10** | EXAM STYLE

Sodium forms Na^+ ions and oxygen forms O^{2-} ions.

(a) Deduce the formula of sodium oxide.

[1 mark]

(b) Draw a complete electron-origin diagram for sodium oxide. Show all ions, brackets, charges and outer-shell electrons.

DIAGRAM SPACE

[6 marks]

**QUESTION 11** | **EXAM STYLE**

Carbon dioxide, CO_2 , contains two carbon-oxygen double bonds.

(a) State the total number of shared electron pairs in one CO_2 molecule.

[1 mark]

(b) Draw a complete dot-and-cross diagram for CO_2 . Show electron origin and all lone pairs.

DIAGRAM SPACE

[6 marks]

**QUESTION 12** | EXAM STYLE

A water molecule reacts with H^+ to form H_3O^+ .

(a) Identify the electron-pair donor and acceptor and state the direction of the dative-bond arrow.

[3 marks]

(b) Draw the H_3O^+ ion, showing the origin of the electron pair in the dative bond, brackets and charge.

DIAGRAM SPACE

[3 marks]

QUESTION 13 | EXAM STYLE

Average bond enthalpy data are shown: $\text{C}-\text{C}$ 347 kJ mol^{-1} ; $\text{H}-\text{H}$ 436 kJ mol^{-1} ; $\text{C}=\text{C}$ 612 kJ mol^{-1} ; $\text{N}\equiv\text{N}$ 945 kJ mol^{-1} .

(a) Identify the strongest bond and explain your choice.

[2 marks]

(b) Use the data to compare the strengths of a $\text{C}-\text{C}$ bond and a $\text{C}=\text{C}$ bond.

[2 marks]

**QUESTION 14** | EXAM STYLE

Magnesium has a giant metallic lattice.

(a) Describe the particles and attraction in metallic magnesium.

.....

.....

.....

[2 marks]

(b) Explain why magnesium conducts electricity as both a solid and a liquid.

.....

.....

.....

[2 marks]

(c) Explain why magnesium has a relatively high melting point.

.....

.....

.....

[2 marks]

QUESTION 15 | EXAM STYLE

Diamond and graphite are giant covalent forms of carbon.

(a) Explain why diamond has a very high melting point.

.....

.....

.....

[2 marks]

(b) Explain why diamond does not conduct electricity.

.....

.....

.....

[2 marks]

(c) Explain why graphite conducts electricity.

.....

.....

.....

[2 marks]

(d) Explain why the layers in graphite can slide over one another.

.....

.....

.....

[2 marks]

**QUESTION 16** | EXAM STYLE

Iodine is a simple molecular substance made from I_2 molecules.

(a) Describe the particles and attractions in solid iodine.

[2 marks]

(b) Explain why iodine has a much lower melting point than diamond.

[2 marks]

(c) State what is overcome when iodine melts and what remains intact.

[1 mark]

(d) Explain why iodine does not conduct electricity in the solid or liquid state.

[1 mark]

QUESTION 17 | EXAM STYLE

The properties of four substances are shown.

Substance	Melting point / °C	Conducts solid	Conducts liquid
A	801	no	yes
B	650	yes	yes
C	3550	no	no
D	114	no	no

(a) Identify the most likely structure type for A, B, C and D.

[4 marks]

(b) For each substance, use one conductivity clue to justify the identification.

[4 marks]

**QUESTION 18** | **EXAM STYLE**

Each statement contains a chemistry error. Identify the error and write a corrected statement.

(a) The ionic bond is the transfer of electrons from a metal atom to a non-metal atom.

.....

.....

.....

.....

.....

[2 marks]

(b) Molten sodium chloride conducts because its electrons become free to move.

.....

.....

.....

.....

.....

[2 marks]

(c) Iodine melts because the covalent bond inside each I_2 molecule is broken.

.....

.....

.....

.....

.....

[2 marks]

(d) All ionic compounds dissolve in water simply because their ions are charged.

.....

.....

.....

.....

.....

[2 marks]



SECTION C

HARDER MIXED

Five less-scaffolded applications in unfamiliar contexts

Independent thinking: Choose the structure-to-property route yourself and keep bond formation separate from the attraction that defines the bond.

QUESTION 19 | HARDER MIXED

A student compares sodium chloride, magnesium and iodine.

(a) State the structure type of each substance.

.....

.....

.....

[3 marks]

(b) Predict whether each conducts electricity as a solid and justify each prediction using the available mobile or fixed particles.

.....

.....

.....

.....

.....

.....

.....

[3 marks]

(c) Explain why sodium chloride has a much higher melting point than iodine.

.....

.....

.....

.....

.....

.....

.....

[2 marks]

**QUESTION 20 | HARDER MIXED**

Element X forms X^{3+} ions. Element Y forms Y^{2-} ions.

(a) Deduce the formula of the compound formed.

[2 marks]

(b) State the total number of electrons transferred when one formula unit forms.

[2 marks]

(c) Draw a complete electron-origin diagram for one formula unit. Show the smallest ion ratio, complete outer shells, brackets and charges.

DIAGRAM SPACE

[6 marks]

**QUESTION 21 | HARDER MIXED**

Four unfamiliar substances have the following properties.

Substance		Melting point / °C	Solid conductivity	Liquid conductivity
P	80		none	none
Q	1414		none	none
R	770		none	good
S	1085		good	good

(a) Identify the most likely structure type for P, Q, R and S.

.....

.....

.....

.....

[4 marks]

(b) Explain the change in conductivity when R melts.

.....

.....

.....

.....

.....

[3 marks]

(c) Explain why Q has a much higher melting point than P.

.....

.....

.....

.....

.....

[2 marks]



QUESTION 22 | HARDER MIXED

Graphite is used as an electrode. Diamond is used on cutting tools.

(a) Explain why graphite is suitable for use as an electrode.

.....

.....

.....

.....

.....

[3 marks]

(b) Explain why diamond is suitable for use on a cutting tool.

.....

.....

.....

.....

.....

[3 marks]

(c) Explain why graphite is softer than diamond.

.....

.....

.....

.....

.....

[2 marks]



A student writes calcium nitrate as CaNO_3 and ammonium sulfate as NH_4SO_4 .

The same student draws the diagram below for $\text{NH}_3 + \text{H}^+ \rightarrow \text{NH}_4^+$.

$$\text{NH}_3 + \text{H}^+ \longrightarrow \text{NH}_4^+$$

(c) Draw a corrected model diagram.

Bonding I | OCR A | 18



SECTION D

STRETCH / SYNOPTIC

One final challenge connecting electron configurations with Bonding I

QUESTION 24 | STRETCH / SYNOPTIC

Stretch note: This question connects Atomic Structure II electron configurations with ionic formulae, electron origin and conductivity.

Atom X has electron configuration $1s^2 2s^2 2p^6 3s^2$. Atom Y has electron configuration $1s^2 2s^2 2p^5$.

(a) Identify X and Y.

[2 marks]

(b) State the ions formed by X and Y.

[2 marks]

(c) Deduce the formula of the compound formed.

[1 mark]

(d) Draw a complete electron-origin diagram for the compound.

DIAGRAM SPACE

[5 marks]

(e) Explain why the compound does not conduct as a solid but does conduct when molten.

[2 marks]

✓ **DONE:** Mark the pack using the worked answers. For every lost mark, identify whether the error came from the definition, electron origin, charge balance or structure-property chain.



CrackingCHEM

GCSE | A-LEVEL

UNDERSTAND • APPLY • SUCCEED

OCR A H432

Bonding I

Fully worked answers | OCR A-style marking guidance

FULLY WORKED. PRECISELY MARKED. BUILT FOR IMPROVEMENT.

OCR A SPECIFICATION

2.1.2(a) | 2.2.2(a-f,n-o) | linked 3.1.1(d-f)

TEST IT	EXAM STYLE	HARDER MIXED	STRETCH
8	10	5	1

24 QUESTIONS MARKED

166 MARKS EXPLAINED

FINAL NOTES-ALIGNED EDITION | FORMULAE • BONDING • STRUCTURE • PROPERTIES

Independent practice material. Every question is original and stays within the finished OCR A Bonding I notes.



OCR A MAP

How this pack matches the finished notes

Full coverage of Bonding I with Bonding II and later Periodicity held back

Specification area	Recurring OCR A demand	Frequent trap	Questions
Ionic formulae	Recall charges; balance charge; use brackets and smallest ratios.	Leaving an unsimplified ratio or omitting brackets around repeated compound ions.	1-2, 10, 20, 23-24
Ionic bonding	Separate electron transfer from electrostatic attraction; prove electron origin.	Calling transfer the bond or omitting brackets, charges or electron totals.	3-4, 9-10, 18-20, 24
Covalent and dative	Use nuclei/shared-pair wording; draw single, multiple and donated pairs.	Incomplete definitions, missing lone pairs or arrows pointing from acceptor to donor.	5-7, 11-13, 23
Metallic and giant covalent	Name particles, attraction, delocalised electrons and structure-property chains.	Using free electrons, ions in graphite, or structure labels without reasoning.	8, 14-15, 17, 19, 21-22
Simple molecular	Distinguish intermolecular forces from covalent bonds during state changes.	Claiming molecules contain ionic bonds or covalent bonds break on melting.	16-19, 21

Research basis: Checked against the final CrackingChem OCR A Bonding I notes, the CrackingChem Year 12 master specification map and OCR A Chemistry H432 specification Version 3.1. Every question is original.

Assessment boundary: No shapes, bond angles, electronegativity, polarity or detailed intermolecular-force types are assessed. Silicon and graphene are not used beyond the limited foundation statement in the notes.

HOW TO SELF-MARK EACH ANSWER

Step	Action
1 Compare	Read the complete model answer before counting marks.
2 Tick	Award one tick only for each distinct marking point shown in brackets.
3 Diagnose	Label the first lost link: definition, charge, electron origin, attraction, energy or mobility.
4 Improve	Rewrite the full answer or redraw the complete diagram without looking.

Diagram marking checklist: Count every outer-shell electron; distinguish its origin; place ionic products in square brackets; write every charge; and make each dative arrow start at the donated pair and point to the acceptor.



SECTION A

TEST IT answers

Direct retrieval with exact OCR A terminology and fully correct diagrams

TEST IT 1

- NO_3^- nitrate [1]; CO_3^{2-} carbonate [1]; SO_4^{2-} sulfate [1]; OH^- hydroxide [1]; NH_4^+ ammonium [1]; Zn^{2+} zinc [1]; Ag^+ silver [1].

Examiner guidance: A correct ion requires both the formula and the charge. A charge written as a subscript is not acceptable.

TEST IT 2


- Aluminium oxide Al_2O_3 [1]; calcium nitrate $\text{Ca}(\text{NO}_3)_2$ [1]; ammonium sulfate $(\text{NH}_4)_2\text{SO}_4$ [1].
- Magnesium hydroxide $\text{Mg}(\text{OH})_2$ [1]; zinc carbonate ZnCO_3 [1]; silver nitrate AgNO_3 [1].
- Use brackets only when more than one of a compound ion is needed, so the subscript applies to the whole ion [1].

TEST IT 3

- (a) Ionic bonding is the electrostatic attraction [1] between positive and negative ions [1]. [2]
- (b) Electron transfer is the process that forms the ions [1]; the bond is the attraction between the ions after transfer [1]. [2]

TEST IT 4

- One Mg^{2+} ion and two Cl^- ions, all bracketed and correctly charged [2].
- Each chloride ion has eight outer-shell electrons [1], with one electron on each chloride shown as transferred from Mg [1]. Magnesium has lost two electrons in total [1]. [5]

 MgCl_2 : electron-origin ionic diagram


• = anion's original electron x = transferred electron from the metal

TEST IT 5

- (a) A covalent bond is the strong electrostatic attraction [1] between a shared pair of electrons [1] and the nuclei of the bonded atoms [1]. [3]
- (b) H_2 has one shared pair; O_2 has two; N_2 has three [2]. [2]

TEST IT 6


- HCl : one shared dot-and-cross pair and three lone pairs on Cl [2]. O_2 : two shared pairs and two lone pairs on each O [2]. N_2 : three shared pairs and one lone pair on each N [3]. [7]

**Model electron-origin diagrams**

• = electron from the left atom × = electron from the right atom Lone pairs sit on the shell line.

TEST IT 7

- (a) A dative covalent bond is a shared pair of electrons [1] in which both electrons were supplied by the same atom [1]. [2]
- (b) NH_3 is the electron-pair donor and H^+ is the acceptor [1]; the arrow begins at the N lone pair and points towards H^+ [1]. [2]
- (c) NH_4^+ is enclosed in square brackets with an overall + charge [1]; the new N-H shared pair contains two electrons originating from N [1]. [2]

Dative-bond formation: NH_3 donates a lone pair to H^+ 

new N-H shared pair = •• (both electrons from N)

• = electron from N × = electron from H Arrow starts at the donated pair and points to H^+ .

TEST IT 8

- (a) Metallic bonding is the strong electrostatic attraction [1] between positive metal ions and delocalised electrons [1]. [2]
- (b) Bond Y is stronger because it has the larger average bond enthalpy, so more energy is needed to break it [1]. [1]



SECTION B

EXAM STYLE answers


Complete routes, precise particles and model electron-origin diagrams

EXAM STYLE 9

- (a) NaCl has many strong electrostatic attractions between oppositely charged ions throughout a giant lattice [1]; much energy is required to overcome them [1]. [2]
- (b) In solid NaCl, ions are fixed in position, so there are no mobile charge carriers [1]. When molten, the ions are mobile [1] and carry charge through the liquid [1]. [3]
- (c) In aqueous NaCl, dissolved Na^+ and Cl^- ions are mobile and carry charge [1]. [1]
- (d) Ionic compounds are often, but not always, soluble in water; solubility depends on whether new water-ion attractions compensate for separating the ions in the lattice [1]. [1]

EXAM STYLE 10


- (a) Two Na^+ ions are needed for one O^{2-} ion, so the formula is Na_2O [1]. [1]
- (b) Two bracketed Na^+ ions and one bracketed O^{2-} ion [2]; oxide has eight outer-shell electrons [1], including two electrons shown as transferred from sodium [2]; electron totals and charges are consistent with Na_2O [1]. [6]

 Na_2O : electron-origin ionic diagram

• = anion's original electron x = transferred electron from the metal

EXAM STYLE 11


- (a) Two $\text{C}=\text{O}$ double bonds contain four shared electron pairs in total [1]. [1]
- (b) Carbon is central with two shared pairs to each oxygen [2]; electron origin is shown consistently in all four shared pairs [2]; each oxygen has two lone pairs and carbon has no lone pairs [2]. [6]

Carbon dioxide: two $\text{C}=\text{O}$ double bonds

• = oxygen electron x = carbon electron Shared pairs are in the overlaps; lone pairs sit on the shell line.

EXAM STYLE 12

- (a) H_2O is the donor [1]; H^+ is the acceptor [1]; the arrow starts at an oxygen lone pair and points to H^+ [1]. [3]
- (b) H_3O^+ is bracketed and has an overall + charge [1]; the newly formed O-H shared pair contains two electrons from O [1]; the remaining outer electrons are complete and consistently shown [1]. [3]

**Dative-bond formation: H_2O donates a lone pair to H^+** **EXAM STYLE 13**

- (a) $\text{N}\equiv\text{N}$ is strongest because 945 kJ mol^{-1} is the largest average bond enthalpy [1], so it requires the most energy to break [1]. [2]
- (b) $\text{C}=\text{C}$ is stronger than $\text{C}-\text{C}$ [1] because 612 kJ mol^{-1} is greater than 347 kJ mol^{-1} [1]. [2]

EXAM STYLE 14

- (a) Magnesium contains positive metal ions and delocalised electrons [1] with strong electrostatic attraction between them [1]. [2]
- (b) Delocalised electrons are mobile in both solid and liquid magnesium [1] and carry electrical charge [1]. [2]
- (c) Strong electrostatic attractions act throughout the giant lattice [1], so much energy is needed to overcome them [1]. [2]

EXAM STYLE 15

- (a) Diamond is a giant three-dimensional covalent network [1]; many strong covalent bonds must be broken, requiring much energy [1]. [2]
- (b) All four outer electrons of each carbon are localised in covalent bonds [1], so there are no mobile charged particles [1]. [2]
- (c) Each carbon in graphite forms three covalent bonds, leaving delocalised electrons [1] that move along the layers and carry charge [1]. [2]
- (d) Only weak attractions act between graphite layers [1], so little energy is needed for layers to slide [1]. [2]

EXAM STYLE 16

- (a) Solid iodine contains neutral I_2 molecules [1] attracted to one another by intermolecular forces [1]. [2]
- (b) Iodine requires only the intermolecular forces between molecules to be overcome [1], whereas diamond requires many strong covalent bonds to be broken [1]. [2]
- (c) Intermolecular forces are overcome; the $\text{I}-\text{I}$ covalent bonds remain intact [1]. [1]
- (d) Neutral molecules provide no mobile ions or delocalised electrons [1]. [1]

EXAM STYLE 17

- (a) A ionic [1]; B metallic [1]; C giant covalent [1]; D simple molecular [1]. [4]
- (b) A conducts only when its ions become mobile in the liquid [1]. B conducts in both states because delocalised electrons remain mobile [1]. C has no mobile charged particles [1]. D contains neutral molecules and no mobile charge carriers [1]. [4]

EXAM STYLE 18

- (a) Transfer forms the ions [1]; ionic bonding is the electrostatic attraction between oppositely charged ions [1]. [2]
- (b) The charge carriers are mobile Na^+ and Cl^- ions [1], not free electrons [1]. [2]
- (c) Melting iodine overcomes intermolecular forces between I_2 molecules [1]; the $\text{I}-\text{I}$ covalent bonds remain intact [1]. [2]
- (d) Ionic compounds are often but not always water-soluble [1]; solubility depends on the balance between attractions broken and new solvent-ion attractions formed [1]. [2]



SECTION C

HARDER MIXED answers


Less-scaffolded comparisons, unfamiliar data and corrected misconceptions

HARDER MIXED 19

- (a) Sodium chloride: giant ionic [1]; magnesium: giant metallic [1]; iodine: simple molecular [1]. [3]
- (b) Solid NaCl does not conduct because its ions are fixed [1]. Solid Mg conducts because delocalised electrons are mobile [1]. Solid iodine does not conduct because neutral I_2 molecules provide no mobile charge carriers [1]. [3]
- (c) NaCl has strong electrostatic attractions throughout a giant lattice [1], whereas melting iodine overcomes weaker intermolecular forces between molecules [1]. [2]

HARDER MIXED 20

- (a) Charge balance requires $2 \times (+3)$ and $3 \times (-2)$, so the formula is X_2Y_3 [2]. [2]
- (b) Two X atoms each lose three electrons, so six electrons are transferred in total [2]. [2]
- (c) Two bracketed X^{3+} ions and three bracketed Y^{2-} ions [2]; each Y^{2-} has eight outer-shell electrons [1], with two transferred electrons shown on each [2]; all charges and the 2:3 ratio are correct [1]. [6]

 X_2Y_3 : electron-origin ionic diagram

• = anion's original electron x = transferred electron from the metal

HARDER MIXED 21


- (a) P simple molecular [1]; Q giant covalent [1]; R ionic [1]; S metallic [1]. [4]
- (b) In solid R, ions are fixed and cannot carry charge [1]. On melting, ions become mobile [1] and carry charge through the liquid [1]. [3]
- (c) Q requires many strong covalent bonds in a giant network to be broken [1], whereas P requires only intermolecular forces between molecules to be overcome [1]. [2]

HARDER MIXED 22


- (a) Graphite contains delocalised electrons [1] that are mobile along the layers [1] and carry electrical charge through the electrode [1]. [3]
- (b) Diamond is a rigid three-dimensional network [1] in which each carbon forms four strong covalent bonds [1], making it very hard and resistant to deformation [1]. [3]
- (c) Graphite layers are held together by weak attractions [1], allowing the layers to slide over one another [1]. [2]

HARDER MIXED 23

- (a) Calcium nitrate is $Ca(NO_3)_2$: one Ca^{2+} requires two NO_3^- ions [2]. Ammonium sulfate is $(NH_4)_2SO_4$: two NH_4^+ ions balance one SO_4^{2-} ion [2]. [4]
- (b) The arrow is reversed; it must begin at the nitrogen lone pair and point to H^+ [1]. The product must be enclosed in square brackets with an overall + charge [1]. [2]
- (c) Correct model: NH_3 donates its lone pair to H^+ and forms bracketed NH_4^+ [2]. [2]

**Dative-bond formation: NH_3 donates a lone pair to H^+** **STRETCH / SYNOPTIC 24**

- (a) X has 12 electrons, so X is magnesium, Mg [1]. Y has 9 electrons, so Y is fluorine, F [1]. [2]
- (b) Mg loses two electrons to form Mg^{2+} [1]; F gains one electron to form F^- [1]. [2]
- (c) One Mg^{2+} requires two F^- ions, so the formula is MgF_2 [1]. [1]
- (d) One bracketed Mg^{2+} and two bracketed F^- ions [2]; each fluoride has eight outer-shell electrons [1], including one electron shown as transferred from Mg [1]; final electron totals and charges are correct [1]. [5]

 MgF_2 : electron-origin ionic diagram

• = anion's original electron x = transferred electron from the metal

- (e) In solid MgF_2 , ions are fixed and cannot carry charge [1]. When molten, the ions are mobile and carry charge [1]. [2]

Final check: Every complete property explanation names the structure, particles and attraction before discussing energy or mobility. Correct the earliest wrong link before reworking the rest.